

# Descriptions of two new species of *Nepenthes* (Nepenthaceae) from Thailand and their phylogenetic analysis based on AFLP technique species confirmation

# Sunya Nuanlaong<sup>1</sup>, Maruay Mekanawakul<sup>2</sup> & Potjamarn Suraninpong<sup>1</sup>

**Summary.** Two new species of *Nepenthes* (Nepenthaceae) from Southern Thailand, *N. bracteosa* sp. nov. and *N. hirtella* sp. nov. are described and illustrated. Descriptions of the species are given compared to their related species, *N. krabiensis* and *N. kerrii*, collected in Thailand. They are separated from *N. krabiensis* and *N. kerrii* by characteristics of leaf shape, pitcher shape, indumentum, and bracteole. Confirmation of these two new species was determined by phylogenetic relationships using the Amplified Fragment Length Polymorphism (AFLP) technique.

Key Words. monkey cup, phylogenetic relationship, pitcher plant, taxonomy.

#### Introduction

The genus Nepenthes, known as the tropical pitcher plant or monkey cup, is in the family Nepenthaceae (order Caryophyllales). The first encounter with Nepenthes in the mid-17<sup>th</sup> century focused on its taxonomy, diversity and distribution. The Nepenthes flora of Indochina and the Philippines, in particular, have attracted much recent attention (McPherson 2009; McPherson 2012). Nepenthes comprises approximately 160 species, with the greatest diversity with many endemic species, in the Philippines and Borneo. Several taxa have been recently described from Indochina: N. bokorensis Mey (Mey 2009), N. thai Cheek (Cheek & Jebb 2009), N. holdenii Mey (Mey et al. 2010), N. mirabilis var. globosa M.Catal. (Catalano 2010a), N. suratensis M.Catal. (Catalano 2010b), N. andamana M.Catal. (Catalano 2010c), N. kerrii M.Catal. & Kruetr. (Catalano 2010d), N. chang M.Catal. (Catalano 2010e), N. rosea M.Catal. & Kruetr. (Catalano 2014), N. kongkandana M.Catal. & Kruetr. (Catalano 2015), N. krabiensis Nuanlaong, Onsanit, Chusangr. & Sura. (Nuanlaong et al. 2016), N. malayensis A.Amin, M.N.Faizal & Dome (Tamizi et al. 2020), N. latiffiana M.N.Faizal, A.Amin & Dome and N. domei M.N.Faizal, A.Amin & A.Latif (Ghazalli et al. 2020). In Thailand, 14 species and one variety of Nepenthes are currently recognised. Of these, nine species including our new species are considered to be closely related and they have been attributed to section *Pyrophytae* Cheek & Jebb (Cheek & Jebb 2016). Seven species in section Pyrophytae are found on the summit areas of mountains, namely, N. chang (300 m a.s.l.), N. kerrii (500 m a.s.l.), N. krabiensis (500 - 600 m a.s.l.), N. rosea (450 -520 m a.s.l.), N. smilesii Hemsl. (16 – 1,500 m a.s.l.), N. bracteosa sp. nov. (710 - 760 m a.s.l.) and N. hirtella sp. nov. (348 - 355 m a.s.l.). While four species, N. andamana, N. kampotiana Lecomte, N. kongkandana and N. suratensis are found at sea level. Generally, plants in section Pyrophytae have tuberous rootstocks, which is an adaptation for fire, and are found in seasonal habitats. Members of section Pyrophytae lack a climbing stem and the upper pitchers (which arise after the inflorescence is first produced) are borne on stems with short internodes. The tendril of the upper pitchers lacks a coil, and the partial-peduncles are 1flowered, with short pedicels/partial-peduncles (Cheek & Jebb 2016). During the dry season, Nepenthes decrease leaf, flower, and pitcher production. The tuberous rootstock is likely to act as a water and nutrient store for survival during periods of unsuitable environmental conditions (Mey et al. 2010).

However, the pyrophytic species are found to be closely related to section *Montanae* Danser (Danser 1928), which share most of their characteristics. Plants from section *Montanae* lack fleshy tubers, have climbing stems, coiled tendrils, and longer pedicels (Cheek & Jebb 2009). In southern Thailand near the Malaysian border, two species namely, *N. sanguinea* Lindl. and *N. thai* Cheek, have been found, which are relatives of Penisular *Montanae*, such as *N. benstonei* C.Clarke, *N. gracillima* Ridl., *N. macfarlanei* Hemsl. and *N. ramispina* Ridl. in highland Malaysia (Cheek & Jebb 2009). However, the two new species are similar in overall appearance and share many characteristics

Accepted for publication 10 August 2021.

<sup>&</sup>lt;sup>1</sup> School of Agricultural Technology, Walailak University, Nakhon Si Thammarat, 80160, Thailand. e-mail: potjamas@hotmail.com

<sup>&</sup>lt;sup>2</sup> School of Science, Walailak University, Nakhon Si Thammarat, 80160, Thailand.

with pyrophytic species such as: long racemose inflorescence, flowering at the rosette stage, seed with reduced filiform appendages, coriaceous narrow leaves, decurrent leaf attachment, tuberous rootstock, and pyrophytic habitat. As a result, the two new species could not be classified as members of section *Montanae*.

Despite previous research, evolution within the genus Nepenthes is still poorly understood. Currently, research in Nepenthes has been focused on their actual diversity. These include biogeographical and environmental processes, and animal-plant interactions that are facilitated mostly by characteristics of pitchers and flowers (Ellison & Adamec 2018). Nepenthes is a crosspollinated plant, in which the wide differential architectures of the phenotypic traits differ resulting from both genetic and environmental factors found in their habitat (Rowe & Speck 2005). Genetic analyses have revolutionised systematics and provided important insights into the understanding plant evolution. The complete genome of Nepenthes has not yet been published, resulting in a limited number of molecular markers available for characterisation of Nephethes taxa. AFLPs, a PCR-based method that selectively amplifies a subset of digested DNA fragments to produce distinctive fingerprints, is the common molecular marker used for comparing genetic differences (Paun & Schönswetter 2012). The effectiveness of AFLP analysis is derived from its capacity to rapidly produce large numbers of marker fragments for any

organism without the requirement of understanding any genomic sequence. The AFLP technique has been used for various applications, including genetic mapping, genealogical studies among closely related individuals, quantification of genetic diversity within and among species, and phylogenetic studies of closely related species (Savelkoul *et al.* 1999).

#### **Materials and Methods**

Morphological analysis

The extent of occurrence (EOO) and area of occurrence (AOO) of the *Nepenthes* populations was calculated to determine the conservation status using Geocat (Bachman *et al.* 2011). All data and dimensions were either collected in the field from live plant specimens, or by a more thorough examination using a Nikon SMZ-U Stereo Zoom Microscope with a Nikon DXM1200F digital camera and ACT-1 software (version 2.70). Morphological terminology applied for the descriptive studies follows Cheek & Jebb (2009) and Nuanlaong *et al.* (2016).

# AFLP analysis and phylogenetic relationships

To determine the putative phylogenetic relationships of the new *Nepenthes* species studied, they were compared with five *Nepenthes* species distributed on mountain tops, specifically *N. chang*, *N. kerrii*, *N. krabiensis*, *N. rosea*, and *N. smilesii*. In addition, *N. thorelii* and *N. mirabilis* were used as an ingroup and an outgroup, respectively (Table 1). Each species included five replications (random plants and sex).

Total genomic DNA was isolated from fresh pitcher tissues using the methods of Suraninpong et al. (2015). Before analysis, DNA from five replicates was pooled in equal quantities. The AFLP analysis was conducted as described by Vos et al. (1995) with some modifications. Initial/first genomic DNA (300 ng) was digested with 10 units of EcoRI (Roche, USA) and 5 units of Tru9I (Roche, USA) in 1X SuRE/Cut™ Buffer A (Roche, USA) to a final volume of 30 µl for 24 h at 37°C. The mixture was added to 10 µl of ligation mix containing 50 pmol adapters of EcoRI and Tru9I, 1 unit of T4-DNA ligase (Roche, USA), and 1x ligation buffer. The ligation reaction was performed at 37°C for 12 h. Then, 5X DNA template (digestion-ligation mixture) dilution was prepared for pre-amplification (PCR I) by adding 2 µl 5X DNA template dilution with 0.25 µM of primer (1 selective base of EcoRI and Tru9I), 1X TopTaq Master Mix Kit (QIAGEN, USA) and dH<sub>2</sub>O, in a final volume of 10 µl. The thermal conditions for PCR I were 20 cycles of 30 s at 95°C, 1 min at 56°C, 1 min at 72°C with a final extension of 7 min at 72°C (T100 Thermal Cycler Bio Red, USA). Then, the PCR I product was diluted to 10X (1 µl PCR I plus with 9 µl dH<sub>2</sub>O) and was mixed with 0.25 µM of primer (2 or 3 selective bases of EcoRI and Tru9I), 1X TopTaq Master Mix Kit and dH<sub>2</sub>O to a final volume of 10 µl. The PCR II was conducted with the annealing temperature reduced by 1°C every cycle. It started with

Table 1. Plant material examined in this study.

#### Species

Nepenthes	mirabilis	(Lour.)	Druce
Nehenthes	thorelij L	ecomte	

Nepenthes chang M.Catal.

Nepenthes kerrii M.Catal. & Kruetr.

Nepenthes smilesii Hemsl.

Nepenthes krabiensis Nuanlaong, Onsanit, Chusangr. & Suran.

Nepenthes rosea M.Catal. & Kruetr.

Nepenthes bracteosa Suran. & Nuanlaong sp. nov.

Nepenthes hirtella Nuanlaong & Suran. sp. nov.

10 cycles of 30 s at 95°C, then 30 s at 65°C, and 1 min at 72°C. Continuously, 30 cycles of 30 s at 95°C, 30 s at 56°C, 1 min at 72°C, 7 min at 72°C and 4°C until the reaction was complete. Then, the reaction was stopped by adding 10 µl of stop-loading dye (10 mM EDTA pH 8.0, 98% formaldehyde, Bromophenol Blue & Xylenecyanol). The PCR product was separated on 5% (w/v) polyacrylamide gel electrophoresis. The DNA bands were visualised via silver staining and were scored manually.

The NTSYSpc (version 2.20j N) program was used for cluster analysis based on a similarity matrix. Genetic distances were obtained using the NTSYS-pc 2.20j statistic package program according to Jaccard's similarity coefficient. The matrix was analysed by the unweighted pair-group method with arithmetic mean (UPGMA) (Rohlf 1998). The relationships between species are illustrated as a dendrogram. The presence and absence of bands were scored as 1 and 0, respectively. Bands of similar size and intensity were assumed to be homologous. The polymorphism (P)was calculated as  $P = (K/N) \times 100\%$ ; where K is the number of polymorphic bands and N is the total number of amplified bands. The polymorphism information content (PIC) value for each AFLP primer combination was calculated based on Roldàn-Ruiz et al. (2000) as PIC<sub>i</sub> =  $2f_i(1 - f_i)$ ; where PIC<sub>i</sub> is the PIC of marker *i*,  $f_i$  is the frequency of the marker bands which were present, and 1 -  $f_i$  is the frequency of marker bands which were absent. The final PIC value was the average of the PIC values of each marker over the bands for each primer combination. The Jaccard's similarity coefficient was computed for each pair of species (Jaccard 1908). The FreeTree program (Pavlicek et al. 1999) was used for the bootstrap analysis with 1,000 resampled datasets. We applied the same levels of support to the Bootstrap values as Richardson *et al.* (2000), which indicates that a value in the range of 85 - 100% means high reliability, 71 -84% means medium reliability, and 50 - 70% means low reliability.

#### **Results and Discussion**

#### Morphological evidence

Several morphological characters allow the new species, namely *Nepenthes bracteosa* and *N. hirtella*, to be distinguished from presently recognised species. The morphology of the two new species differs distinctly from *N. krabiensis* and *N. kerrii* as shown in Table 2.

#### AFLP and phylogenetic relationship

In this study, more than 50 primer combinations from EcoRI and Tru9I were examined, but only eleven primer combinations were clearly classified for polymorphic profiling on nine Nepenthes species (Table 3). A total of 1,415 DNA fragments scored across all primer sets were observed, out of which 1,323 (93.30%) were polymorphic, which enabled differentiation of the accessions analysed. The size of AFLP fragments generated by different primer combinations ranged from 200 to 800 bp. The highest number of polymorphisms were for ER-AC/Tru-CGT (98.64%) followed by ER-AC/Tru-CAG (94.96%) and ER-AGC/ Tru- CAG (94.44%). Conversely, the lowest polymorphism at 90.65% was found on ER-AT/Tru -CAG primer combinations. Monomorphism was lower than polymorphism. Low monomorphism indicates that there is a high degree of variation between the species. The PIC value ranged from 0.71 to 0.86 with an average of 0.76 per fragment. The highest value 0.86 was obtained for the ER-AC/Tru-CGT primer combination, followed by 0.78 for ER-ATG/Tru-GAG. A strong correlation between polymorphism and PIC  $(r^2=0.86)$  was revealed.

Jaccard's similarity coefficient was found to vary from 0.22 to 0.65, with an average of 0.42, suggesting a high genetic diversity among the nine *Nepenthes* species (Table 4). The dendrogram from the cluster analysis, constructed from 11 AFLP markers, indicates that the *Nepenthes* species can be divided into two groups and two separate individual species at 0.46 of similarity coefficient with a bootstrap value of 100% (Fig. 1). This grouping was consistent with the existing morphological and geographical classification of *Nepenthes* species and was similar to previous reports of the relationships from geographical and genetic distribution by Suraninpong *et al.* (2015).

From the results, the first group contains two subgroups, the first of which is Nepenthes rosea, and the second subgroup is N. krabiensis, N. bracteosa, N. kerrii and N. hirtella. All species of this group were distributed throughout southern Thailand but confined to different places. Nepenthes bracteosa clearly showed separation from N. krabiensis with a similarity coefficient of 0.65 with a bootstrap value of 88%. Nebenthes hirtella was identified in the same related subgroup with N. krabiensis and N. bracteosa. From the result of the low similarity coefficient (0.45 - 0.65), high genetic variation among these species was discovered. In contrast, the species from North-East Thailand namely, N. chang and N. smilesii were separated into the second subgroup and were shown to have a similarity coefficient of 0.50 with a bootstrap value of 96%. The results published here suggest that all Nepenthes species found in Thailand are derived from a common ancestor, of which N. thorelii Lecomte was the first species separated from the aggregate group. Likewise, the phylogenetic relationship infor-

**Table 2.** Comparisons of geography and selected plant characteristics of Nepenthes bracteosa and N. hirtella with N. krabiensis and N. kerrii.

	N. krabiensis	N. bracteosa	N. kerrii	N. hirtella
Geography	SW Thailand, 500 – 600 m a.s.l.	SE Thailand, 710 – 760 m a.s.l.	Island, SW Thailand, 500 m a.s.l.	SW Thailand, 348 – 355 m a.s.l.
Leaf shape	lanceolate	linear to lanceolate	obovate	oblanceolate
Leaf apex	acute	acute to acuminate	acuminate	acuminate
Lid of lower pitcher	broadly ovate	broadly ovate to round	round	ovate
Spur	filiform	filiform with branches	simple or branches	filiform with branches
Male inflorescence	2-flowered, partial peduncles	1-flowered	1-flowered	1-flowered or rarely 2- flowered, partial peduncles
Androphore length (mm)	4	2.0 – 2.5	1.5	1.0 - 2.0
Bracts	male inflorescence: present at base or lower half of the pedicel female inflorescence:	present all flowers of both male and female inflorescences	absent	present at base of rachis (1 – 2 flowers) both male and female inflorescences
	absent			
Indumentum	lacking on stem, lamina, and leaf margin	lacking on stem, lamina, leaf margin, and shoot	covering leaf axil and inflorescence	covering all vegetative parts

Primer pairs	Total no. of bands	PB	MB	Р	M	PIC
ER-AG/Tru-CAG	100	92	8	92.00	8.00	0.74
ER-AG/Tru-CGT	101	93	8	92.08	7.92	0.71
ER-AT/Tru-CAG	107	97	10	90.65	9.35	0.71
ER-AC/Tru-CAG	139	132	7	94.96	5.04	0.74
ER-AC/Tru-CGT	147	145	2	98.64	1.36	0.86
ER-AGA/Tru-CGA	113	106	7	93.81	6.19	0.76
ER-AGA/Tru-CCA	136	127	9	93.38	6.62	0.76
ER-AGA/Tru-CAG	119	108	11	90.76	9.24	0.72
ER-AAG/Tru-CAG	149	139	10	93.29	6.71	0.75
ER-AGC/Tru-CAG	162	153	9	94.44	5.56	0.79
ER-ATG/Tru-GAG	142	131	11	92.25	7.75	0.78
Total	1415	1323	92	1026.26	73.74	8.32
Mean	128.64	120.27	8.36	93.30	6.70	0.76

**Table 3.** Description of AFLP results from seven *Nepenthes* species and two new species, *N. bracteosa* and *N. hirtella* in Thailand: PB polymorphic bands; **MB** monomorphic bands; *P* percentage of polymorphism; *M* percentage of monomorphism; **PIC** polymorphism information content.

mation explains the genetic link between the two newly discovered *Nepenthes* species, *N. bracteosa* and *N. hirtella*. *Nepenthes bracteosa* and *N. hirtella* are separated from *N. krabiensis* by their genetic background, even though they are found in close proximity.

#### **Taxonomic Treatment**

**1. Nepenthes bracteosa** *Suran. & Nuanlaong* **sp. nov.** Type: Thailand, Nakhon Si Thammarat Province,

August 2019, *Nuanlaong* BKF197454 (holotype BKF!; isotype PSU!).

Scandent, glossy climbing herb. *Root* branched, fleshy tubers. *Stem* terete, 0.5 - 1.0 m long, 0.3 - 0.5 cm diam.; internodes 1.3 - 2.7 cm long; axillary bud present in 2 - 3 axils in the upper part of the plant when flowering; upper part green to red and lower part brown. *Leaves* pseudo-petiolate, coriaceous, linear

to lanceolate;  $14.0 - 24.0 \times 2.0 - 3.9$  cm, longitudinally folded to form a V-shape; blade apex acute to acuminate, base attenuate, entire, dilating at the node, projecting along the stem as a wing, clasping the stem by three-quarters of its circumference, decurrent for 0.8 – 2.8 cm of its length, obscure longitudinal nerves on each side of the midrib at the edge of lamina, as leaves are thickened, inconspicuous reticulated pinnate venation; midrib conspicuous; tendrils straight, terete, 15.0 - 26.0 cm long in lower pitchers, upper pitchers 10.0 – 15.0 cm long, coiling; glossy light green when young, green when mature, emerging from an orange-amber, triangular lateral bud at the base of the midrib near the leaf attachment. Lower pitcher c.  $10.5 - 19.0 \times 3.4 - 5.6$  cm; ovate in the lower part with digestive glands on the inner surface, cylindrical to narrowing in the upper part; conspicuous midsection hip; with two fringed wings, 0.6 - 1.3 cm wide, along the length of the pitcher, 6.5 - 12.0 cm from pitcher mouth to tendril, fringed elements 0.3 - 1.2 cm long, 4 - 8 fringed elements per cm; mouth ovate, concave; peristome cylindrical, undulate, revolute when pitcher mature, 0.7 - 1.9 cm wide, inner edge with teeth 0.6 - 1.91.7 mm long, ridges approximately 0.4 – 0.5 mm apart;

Table 4. Percentage similarity matrix for nine Nepenthes accessions, generated from Jaccard's similarity coefficient (Jaccard 1908). S1 N. rosea; S2 N. krabiensis; S3 N. bracteosa; S4 N. hirtella; S5 N. smilesii; S6 N. kerrii; S7 N. chang; S8 N. thorelii; S9 N. mirabilis.

	<b>S</b> 1	S2	<b>S</b> 3	<b>S</b> 4	<b>S</b> 5	<b>S</b> 6	<b>S</b> 7	<b>S</b> 8	<b>S</b> 9
<u>S1</u>	1.00								
<b>S2</b>	0.62	1.00							
<b>S</b> 3	0.49	0.65	1.00						
<b>S4</b>	0.45	0.56	0.53	1.00					
<b>S</b> 5	0.42	0.48	0.45	0.43	1.00				
S6	0.48	0.60	0.54	0.52	0.46	1.00			
<b>S</b> 7	0.44	0.51	0.47	0.45	0.50	0.50	1.00		
<b>S</b> 8	0.36	0.39	0.38	0.34	0.35	0.37	0.39	1.00	
S9	0.22	0.24	0.24	0.23	0.24	0.24	0.25	0.22	1.00



Fig. 1. UPGMA-based dendrogram showing genetic relationships among the nine *Nepenthes* species used in this study. The dendrogram is based on the genetic distance calculated according to Jaccard's similarity coefficient with 1000 bootstrap.

lid broadly ovate to round,  $2.8 - 4.4 \times 3.3 - 5.0$  cm, larger than mouth, sometimes vaulted or bent towards the mouth, base cordate 1.0 - 4.0 mm, apex occasionally emarginate, indurate, like appendage on the lower surface, nectar glands numerous, elliptic, 0.16 -0.26 mm diam. along the midrib, and with crateriform glands 0.12 - 0.18 mm diam. dispersed over the lower surface of the lid; spur filiform, 0.2 - 0.8 cm long, with branches; green to red on lower part and red on upper part with red speckled outside, light green to red with red blotches over the inside, absent in digestive zone, peristome green to dark red, lid green to red. Upper pitcher tubulose, with a conspicuous hip one third from the base; 10.0 - 19.0 cm long, 2.0 -3.4 cm wide; two fringed wings, 0.1 - 0.3 cm wide, along the length of the pitcher, 9.0 - 16.0 cm long, fringe elements 0.3 - 0.7 cm long, 3 - 6 per cm; pitcher mouth similar to lower pitcher; peristome cylindrical, revolute when pitcher mature, 0.4 - 1.0 cm wide, inner edge with teeth 0.29 - 0.43 mm long, ridges approximately 0.4 - 0.5 mm apart; lid ovate to round,  $2.1 - 3.4 \times 2.1 - 4.4$  cm, base cordate 1.0 - 3.0mm, apex emarginate similar to lower pitcher, elliptic nectar gland along the midrib on the lower surface of lid, numerous near base, 0.1 - 0.2 mm diam. and with crateriform glands 0.05 - 0.1 mm diam. scattered over the lower surface of the lid; spur filiform, 3.0 - 7.0 mm long, with branches; pitchers green or green to red, sometimes red-speckled outside, light green with red blotches over the inner surface, absent in digestive zone, peristome and lid light green. Male inflorescence a raceme, 70.0 – 85.0 cm long, peduncle 47.0 – 55.0 cm long, 0.2 - 0.5 cm diam., rachis 13.0 - 32.0 cm long, flowers 84 - 135, solitary, pedicels 3.0 - 5.0 mm long, with bracts at the base or lower half of pedicel 0.8 -1.0 mm long, present on all flowers; tepals 4, ovate, apex obtuse,  $4.0 - 5.0 \times 3.0 - 4.0$  mm, reflexed in mature flower, densely covered with orbicular to elliptic nectar glands 0.20 - 0.36 mm diam.; androphore 2.0 – 2.5 mm long, anther head globular 2.1 – 2.5 mm diam., anther basifixed, synandrium, 8 – 13 anther sacs, longitudinal dehiscence; tepals light green when young, red when old. Female inflorescence a raceme, 50.0 - 86.0 cm long, 0.2 - 0.4 cm diam., peduncle 42.0 - 74.0 cm long, rachis 9.0 - 16.0 cm long, flowers 20 - 42, solitary, pedicels 0.5 - 1.7 cm long, with bracts 1.0 - 1.5 mm long, inserted at one third of pedicel height, conspicuous on upper part of rachis, present on all flowers; tepals 4, ovate to elliptic,  $3.8 - 4.0 \times 2.0 - 2.2$  mm, nectar glands similar to male inflorescence, 0.20 - 0.28 mm diam., green when young, green to brown when old; ovary superior, with four syncarpous carpels, oblong, 5.0 - 6.0 mm long, 2.3 - 2.5 mm diam., stigma head 1.0 - 1.5 mm diam. Infructescence similar to female inflorescence; tepals persistent; four valves, 1.0 - 1.2 cm long with septicidal capsule; seeds linear, 6.0 - 6.6 mm long c. 98 - 125 seeds per pod. Indumentum conspicuous, inflorescence and outer pitcher pubescent with simple brown hairs; tendril with simple and branched hairs; wing fringed

elements puberulent, with branched brown hairs; leaf midrib puberulent with inconspicuous, simple, dark green hairs; absent on stem, leaf axils, lamina, leaf margin, and shoot. Figs 2 - 4.

**RECOGNITION.** Nepenthes bracteosa belongs to section Pyrophytae, a section distributed in the strongly seasonal region of Indochina. It is closely related to N. krabiensis, but differs in its shorter stem 0.5 - 1.0 m (vs 1.5 - 2.5 m), spur filiform with branches (vs filiform with no branches), axillary bud present in 2 -3 axils in the upper part of plant when flowering (vs a single axillary bud), fringed wing present in the upper pitcher (vs absent), occasionally emarginate and indurate lid apex (vs a blunt apex and not indurate), solitary flowers both male and female inflorescences (vs 2-flowered partial peduncles in male inflorescence and solitary flower in female inflorescence), bracts present on all flowers of both male and female inflorescences (vs present at the base or lower half of the pedicel in male inflorescence but absent in female inflorescence), short and rophore  $(2.0 - 2.5 \text{ mm vs } 4.0 \text{ m$ mm), and 8 - 13 anther sacs (vs 15 - 22 anther sacs). DISTRIBUTION. Nakhon Si Thammarat Province, Thailand (Map 1).

SPECIMENS EXAMINED. THAILAND.

29 August 2019, *Nuanlaong* BKF197454 (holotype BKF; isotype PSU).

**HABITAT**. On summit areas, in sandy soil, generally growing in open forest and limestone rock crevices, 710 – 760 m a.s.l.

**CONSERVATION STATUS**. This species is distributed at a single site A

decline in the quality of the habitat will lead to the disturbance of the long-term survival of the remaining population and may thus increase the extinction risk of this species. The calculation of EOO as  $0.034 \text{ km}^2$  and AOO as  $4.00 \text{ km}^2$  together with the information above suggests that this species is Critically Endangered (CR) [B1a,b (iii) + B2a,b (iii)] (IUCN 2019).

**PHENOLOGY**. Flowering June to December.

**ETYMOLOGY**. From the Latin "*bracteosa*" derived from "*bracteosus*": bracts are present on the pedicels of all flowers of both male and female inflorescences.

**VERNACULAR NAME.** From the Thai language "Seua-Nakhonsi": "Seua" means the red-speckled tiger-like pattern on the pitcher surface and "Nakhonsi" means *Nepenthes* of Nakhon Si Thammarat Province.

**NOTES.** Nepenthes bracteosa was first discovered from a photograph taken by a tourist. It is only found at a single location in Southern Thailand. It belongs to section *Pyrophytae* on account of sharing several characteristics such as long racemose inflorescence, climbing stem, flower from the rosette stage, tuberous

rootstocks, and pyrophytic habitat. *Nepenthes bracteosa* is similar to *N. krabiensis*, with which it shares being glossy, climbing, its leaf shape and an inconspicuous indumentum, but differs in many other morphological characteristics. However, the results of the phylogenetic relationship revealed a similarity coefficient as 0.65 between *N. bacteosa* and *N. krabiensis*, which is also the highest of all species pairs recorded (Table 3).

**2. Nepenthes hirtella** *Nuanlaong & Suran.* **sp. nov.** Type: Thailand, Krabi Province,

lower pitchers, 29 August 2019, Nuanlaong BKF197453 (holotype BKF!; isotype PSU!).

Scandent, climbing herb. Root branched, fleshy tubers. Stem terete, 0.8 - 1.2 m long, 0.35 - 0.65 cm diam.; internode 1.5 - 5.5 cm long; upper part of the stem green, lower part of the stem brown. Leaves pseudopetioles, coriaceous, oblanceolate; 15.8 - 29.5 × 3.5 -5.4 cm, longitudinally folded to form a V-shape, blade apex acuminate, base attenuate, entire, dilating at the node, projecting along the stem as wing, clasping the stem by three quarters of its circumference, decurrent for 1.7 - 4.3 cm of its length, leaf thickness 0.5 - 1.0mm, inconspicuous longitudinal nerves 3 - 4 on each side of the midrib in outer third of lamina, inconspicuous lamina reticulated pinnate venation; midrib conspicuous; light green in a rosette when young, green when mature, emerging from a red triangular lateral bud at the base of the midrib near the leaf attachment. Lower pitcher c.  $14.0 - 19.7 \times 4.4 - 5.9$  cm; ovate at the lower part with digestive zone at the inner surface, narrowing at the upper part; hip conspicuous above digestive zone; with two fringed wings, 0.3 -0.9 cm wide, along the length of the pitcher, 9.0 -14.3 cm from pitcher mouth to tendril, serrate to double serrate, fringed elements 0.4 - 0.9 cm long, 4 -11 fringed elements per cm; mouth oblique ovate, concave; peristome cylindrical, entire to repand, revolute when pitcher mature, 0.9 - 1.5 cm wide, inner edge with teeth 0.64 - 1.48 mm long, ridges approximately 0.72 - 1.2 mm apart; lid ovate, 3.9 - 4.9  $\times$  3.3 – 4.9 cm, repand, base cordate 0.5 – 3.0 mm; absent appendage, nectar glands numerous, elliptic 0.18 - 0.28 mm diam. along the midrib, with crateriform gland 0.09 - 0.28 mm diam. dispersed over the lower surface of the lid; spur filiform, 0.1 -0.9 cm long, with branches; tendril straight, 12.0 -20.0 cm long, 2.0 – 3.5 mm diam., approximately 10 – 30% longer than pitcher, mostly on the side of the pitcher; light green to yellow or green to red on the lower and green to red or red on the upper part with



**Fig. 2.** Nepenthes bracteosa. A leaf shape and lower pitcher; B upper pitcher; C stem; D male inflorescence; E lower surface of lid with two types of nectar glands and a ridge-like appendage at the lid apex; F male flower; G female flower after pollination; H peristome; J filiform spur with branches; K peristome in transverse section. Drawn from live specimens. DRAWN BY CHAYAN YODTHAMMARAT.



**Fig. 3.** Nepenthes bracteosa. A lower pitcher; B upper pitcher and habitat; C infructescence; D axillary bud present in 2 – 3 axils when flowering; E – G male inflorescence (immature, young, and old flowers); H ridge-like appendage under lid apex; J branched hairs on fringe elements; K nectar glands on lower surface of lid. PHOTOS: POTJAMARN SURANINPONG & SUNYA NUANLAONG.



**Fig. 4.** Nepenthes bracteosa. A lower pitcher; B upper pitcher; C – D habit and habitat. Photos: POTJAMARN SURANINPONG & SUNYA NUANLAONG.

red blotches on the outside surface, light green present or absent red blotches on inside surface, absent in digestive zone, peristome light green, green to red or red peristome, lid green or green to red. Upper pitcher tubulose, with a conspicuous hip one third from the base or middle of pitcher; 13.0 - $21.0 \text{ cm} \log, 0.9 - 2.8 \text{ cm}$  wide; two fringed wings, 1.5 $-8.0 \times 0.1 - 0.2$  cm, which run down the ventral exterior surface from pitcher mouth to the middle or tendril of the pitcher, serrate or double serrate, fringed elements 0.5 - 2.0 mm long, 4 - 8 per cm; mouth oblique ovate; peristome cylindrical, revolute when pitcher mature, 0.3 - 1.0 cm wide, inner edge with teeth 0.48 - 0.84 mm long, ridges approximately 0.92 - 1.40 mm apart; lid ovate,  $2.1 - 4.3 \times 2.0 - 4.3$ cm, base cordate 0.5 - 1.0 mm, nectar gland similar to lower pitcher; spur filiform, 2.0 – 7.0 mm long; tendril straight, 7.0 - 15.0 cm long, 1.0 - 3.0 mm diam., coiling, shorter than pitcher; green or green to orange outside surface with or without red blotches, green with red blotches over inner surface, absent in digestive zone, peristome light green sometimes red stripes present, lid green. Male inflorescence a raceme, 65.0 - 105.0 cm long, 0.3 - 0.4 cm diam., peduncle 44.0 - 56.0 cm long, rachis 22.5 - 47.0 cm long, flowers 68 - 194, solitary flowers or rarely two-flowered partial peduncle, pedicels 0.5 - 0.8 cm long, bracts present at base of peduncle 0.5 - 3.0 mm long, absent on upper half; tepals 4, elliptic to obovate,  $2.0 - 4.0 \times 1.0 - 2.0$ mm, explanate and not reflexed, cruciferous, densely covered with orbicular to elliptic nectar glands 0.35 -0.70 mm diam.; and rophore 1.0 - 2.0 mm long, anther head globular, 1.0 – 2.0 mm diam., anther basifixed, synandrium, 8 - 12 anther sacs, longitudinal dehiscence; tepals light green when young, green to red when old. *Female inflorescence* a raceme, 65.0 - 75.0 cm long, 0.30 - 0.35 mm diam., peduncle 40.0 - 58.5 cm long, rachis 12.0 – 20.0 cm long, bracts similar to male inflorescence; flowers 26 - 57, solitary, pedicels 0.2 -1.0 mm long; tepals 4, elliptic to ovate,  $2.0 - 3.0 \times 1.0 -$ 2.0 mm, nectar glands similar to male inflorescence, 0.35 - 0.60 mm diam.; tepals green when young, green to red when old; ovary superior, with four syncarpous carpels, oblong, 3.0 - 5.0 mm long, 2.0 - 2.5 mm diam., stigma head 1.5 - 2.0 mm diam. Infructescence similar to female inflorescence; tepals persistent; four valves, 1.5 - 1.6 cm long with septicidal capsule; seed linear, 4.0 - 8.0 mm long c. 73 - 109 seeds per pod. Indumentum conspicuous, simple brown hairs, pubescent, 0.17 - 0.30 mm long, present on all vegetative parts; fringed elements puberulent. Figs 5 - 7.

**RECOGNITION.** Nepenthes hirtella belongs to section Pyrophytae, a section distributed in strongly seasonal Indochina. It is closely related to N. kerrii which grows

in Tarutao Marine Park, Satun Province, Thailand. *Nepenthes hirtella* differs from *N. kerrii* in having an oblanceolate leaf shape (vs obovate), tendrils longer than pitcher approximately 10 - 30% (vs longer than pitcher approximately 40 - 50%), tendrils displaced to the side of lower pitcher (vs tendril centrally placed at base of lower pitcher), longer lower pitcher (14.0 – 19.7 cm vs 6.0 – 14.0 cm), lid ovate (vs round), spur filiform (vs simple or branched), solitary flowers or rarely two-flowered partial peduncle (vs solitary flowers), and indumentum covering all the vegetative parts (vs covering leaf axil and inflorescence).

**DISTRIBUTION**. Krabi Province, Thailand (Map 1). **SPECIMENS EXAMINED. THAILAND**.

lower pitchers, 29 August 2019, *Nuanlaong* BKF197453 (holotype BKF!; isotype PSU!). HABITAT. On summit areas, in sandy soil, generally growing in open forest, savannas and scrubland, at 348 – 355 m a.s.l.

**CONSERVATION STATUS.** *Nepenthes hirtella* is found only on a single site

A decline in the quality of the habitat will lead to the disturbance of the long-term survival of the remaining population. It is easy to get to the habitat, which might put this species at risk of extinction. The calculation of EOO as  $0.053 \text{ km}^2$ , AOO as  $8.00 \text{ km}^2$  together with the information above suggests that this species is Critically Endangered (CR) [B1a,b (iii) + B2a,b (iii)] (IUCN 2019).

**PHENOLOGY**. Flowering June to December.

**ETYMOLOGY**. From the Latin "*hirtella*" derived from "*hirtellus*": short hairs are present on all the vegetative parts of the plant and rather hairy.

**VERNACULAR NAME.** From the Thai language "Seua-Amphon": "Seua" means the red-speckled tiger-like pattern on the pitcher surface and "Amphon" is the name of the first collector, Mr Amphon Dumkliang who found them in 2014 – 2015.

NOTES. Nepenthes hirtella was first discovered in 2014 – 2015 by a Nepenthes collector/seller and was known as "Seua-Amphon". Nepenthes hirtella is a member of section Pyrophytae like N. bracteosa and is only found in Krabi Province, Thailand.

Nepenthes hirtella is closely related to N. kerrii but differs in several morphological details, in particular in tendril and indumentum. Nepenthes hirtella is covered by brown hairs throughout, while in N. kerrii the indumentum is confined specifically to the leaf axil and inflorescence. It is distributed on the mainland while N. kerrii is distributed on an island. Moreover, the phylogenetic relationship results show a low similarity index between them (Table 4).



**Fig. 5.** Nepenthes hirtella. A upper pitcher; B lower pitcher with leaf shape; C lower surface of lid with two types of nectar glands; D rear view of pitcher mouth of B; E spur of lower pitcher; F peristome of lower pitcher in transverse section; G female inflorescence after pollination; H male inflorescence; J habit. Drawn from live specimens. DRAWN BY CHAYAN YODTHAMMARAT.



**Fig. 6.** Nepenthes hirtella. A upper pitcher and habitat; B leaves and lower pitcher; C lower pitcher; D male inflorescence; E female inflorescence; F anther head; G peristome with teeth; H branched hairs on fringed element; J nectar glands on lower surface of lid; K stem and leaf axil; L brown hairs on lamina and leaf margin. PHOTOS: POTJAMARN SURANINPONG & SUNYA NUANLAONG.

# Key to Nepenthes species in Thailand

1.	Tuberous rootstock	2
1.	Non-tuberous rootstock	12
2.	Inflorescence racemose with flowers all arranged singly (a 1-flowered) or rarely 2-flowered partial peduncle	e 3
2.	Inflorescence racemose with 2-flowered partial peduncles	10



Fig. 7. Nepenthes hirtella. A lower pitcher; B upper pitcher; C – D habit and habitat. Photos: POTJAMARN SURANINPONG & SUNYA NUANLAONG.

3.	Leaves obovate or oblance late $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots 4$
3.	Leaves linear-lanceolate
4.	Leaves obovate, tendril longer than pitcher approximately $40 - 50\%$ ; indumentum covering leaf axil and
4	Inflorescence
4.	norts
5	Short hairs present on all parts or some part of plant
5. 5	Stem and leaves glabrous (nitcher inflorescence tendril excluded)
6	Indumentum on all vegetative parts: tendril as long as or shorter than pitcher <i>N</i> . smilesi
6.	Indumentum present on some parts of the plant: tendril longer than pitcher
7.	Lower pitcher narrowly ovate in the lower third or lower half and tubular above
7.	Lower pitcher ovate in lower half and narrowing above or completely ovate
8.	Lid smaller than mouth, triangular mouth as large as $\frac{1}{3}$ to $\frac{1}{2}$ of the lower pitcher length; flower green with red
	margins
8.	Lid larger than mouth, ovate mouth as large as 1/4 of the lower pitcher length; flower red N. andamana
9.	Pseudo-petiolate leaves; lid broadly ovate to round; bract present in all flowers in both male and female
	inflorescences
9.	Sub-petiolate leaves; lid orbicular; bract present on partial peduncles
10.	Hip present at the mid-section or absent in lower pitcher; indumentum covering all aerial parts except upper
10	surface of leaves
10.	Hip present at the lower $\frac{1}{3}$ in both lower and upper pitchers; indumentum present on some parts of the
11	plant
11.	numeritum often covering lear margin and midro; lower pitcher green to light pink, with dark speckles
11	Indumentum lacking on stem lamina leaf margin and shoot: lower pitcher green to red with red speckles
11.	outside and light green or green with blotches over the inside
12.	Leaves petiolate: fimbriate leaf margin
12.	Leaves sessile or with a short, poorly defined petiole; entire leaf margin
13.	Lower pitcher ovoid in the basal third, narrowing slightly to form a hip and cylindric above N. mirabilis
13.	Lower pitchers round to oval N. mirabilis var. globosa
14.	Producing ground pitchers
14.	Not producing ground pitchers
15.	Lower pitcher urceolate, peristome incurved; inflorescence paniculate N. ampullaria
15.	Lower pitcher elongated and smaller, very thin peristome; inflorescence racemose N. gracilis
16.	Leaf base amplexicaul, clasping the stem transversely; lid ovate, truncate at base N. sanguinea
16.	Leaf base decurrent, running longitudinally down the stem as wings; lid elliptic, round at base N. thai

#### Acknowledgements

This work was supported by The Plant Genetics Conservation Project under the Royal Initiation of Her Royal Highness Princess Maha Chakri Sirindhorn, Walailak University and Walailak University Fund (Grant No.WU\_IRD61\_11).

We also thank Dr Sarayut Onsanit and Asst. Prof. Dr Suraphon Thitithanakul, Prince of Songkhla University, Suratthani Campus for supporting the fieldwork. Special thanks are given to Mrs Raumporn Ketsarapong, Walailak Botanic Garden for her valuable suggestions and Mr Chayan Yodthammarat for assistance with the illustrations and photographs.

### References

- Bachman, S., Moat, J., Hill, A. W., de la Torre, J. & Scott, B. (2011). Supporting Red List threat assessments with GeoCAT: Geospatial conservation assessment tool. *ZooKeys* 150: 117 – 126.
- Catalano, M. (2010a). Nepenthes mirabilis var. globosa M. Catal. var. nov. In: M. Catalano, Nepenthes della Thailandia: Diario di viaggio. WOW s.r.o., Prague. 40.

(2010b). Nepenthes suratensis M. Catal. sp. nov. In:
 M. Catalano, Nepenthes della Thailandia: Diario di viaggio. WOW s.r.o., Prague. 36.

(2010c). Nepenthes andamana M. Catal. sp. nov. In:
 M. Catalano, Nepenthes della Thailandia: Diario di viaggio. WOW s.r.o., Prague. 34.

\_\_\_\_ (2010d). *Nepenthes kerrii* M. Catal. et T. Kruetr. sp. nov. In: M. Catalano, *Nepenthes* della Thailandia: Diario di viaggio. WOW s.r.o., Prague. 32.

- \_\_\_\_ (2010e). *Nepenthes chang* M. Catal. sp. nov. In: M. Catalano, *Nepenthes* della Thailandia: Diario di viaggio. WOW s.r.o., Prague. 38.
- \_\_\_\_ (2014). *Nepenthes rosea*, una nuova specie dalla Thailandia peninsulare. *AIPC Mag.* 36: 24 – 31.
- (2015). *Nepenthes kongkandana*, da ufficiosa ad ufficiale. *AIPC Mag.* 37: 4 11.
- Cheek, M. & Jebb, M. (2009). Nepenthes group Montanae (Nepenthaceae) in Indo-China, with N. thai and N. bokor described as new. Kew Bull. 64: 319 – 325.
- \_\_\_\_\_ & \_\_\_\_\_ (2016). Nepenthes section Pyrophytae. Planta Carniv. 38 (1): 44 45.
- Danser, B. H. (1928). The Nepenthaceae of the Netherlands Indies. Bull. Jard. Bot. Buitenzorg III. 9: 249 – 438.
- Ellison, A. M. & Adamec, L. (2018). *Carnivorous Plants: Physiology, ecology, and evolution.* Oxford University Press, Oxford.
- Ghazalli, M. N., Tamizi, A. A., Nikong, D., Besi, E. E., Mat-Esa, M. I., Mohd-Nordin, A. R., Latiff, A., Zaini, A. Z. & Shakri, M. A. (2020). *Nepenthes latiffiana* and *N. domei* (Nepenthaceae), two new species of pitcher plants from Terengganu, Peninsular Malaysia. *Webbia.* 75 (1): 5 – 28.
- IUCN Standards and Petitions Committee. (2019). Guidelines for Using the IUCN Red List Categories and Criteria. Version 14. Prepared by the Standards and Petitions Subcommittee, IUCN, Gland. Downloadable from http://www.iucnredlist.org/documents/ RedListGuidelines.pdf. [Accessed 10 Nov. 2021].
- Jaccard, P. (1908). Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaud. Sci. Nat.* 44: 223 – 270.
- Mey, F. S. (2009). Nepenthes bokorensis, a new species of Nepenthaceae from Cambodia. Carniflora Aust. 7 (1): 6 – 15.
- \_\_\_\_, Catalano, M., Clarke, C., Robinson, A., Fleischmann, A. & McPherson, S. (2010). Nepenthes holdenii (Nepenthaceae), a new species of pyrophytic pitcher plant from the Cardamom Mountains of Cambodia. In: S. R. McPherson, Carnivorous Plants and their Habitats. Vol. 2: 1306 – 1331. Redfern Natural History Productions, Poole.
- McPherson, S. (2009). *Pitcher Plants of the Old World 1st ed.* Redfern Natural History, Poole.
- (2012). *New Nepenthes Volume 1, 1st ed.* Redfern Natural History, Poole.
- Nuanlaong, S., Onsanit, S., Chusangrach, V. & Suraninpong, P. (2016). A new species of *Nepenthes*

(Nepenthaceae) from Thailand. *Thai Forest Bull.*, *Bot* 44 (2): 128 – 133.

- Paun, O. & Schönswetter, P. (2012). Amplified Fragment Length Polymorphism (AFLP) — an invaluable fingerprinting technique for genomic, transcriptomic and epigenetic studies. *Methods Molec. Biol.* 862: 75 – 87.
- Pavlicek, A., Hrda, S. & Flegr, J. (1999). Free-Tree Freeware program for construction of phylogenetic trees on the basis of distance data and bootstrap/ jackknife analysis of the tree robustness. Application in the RAPD analysis of the genus *Frenkelia*. *Folia Biol. (Prague)* 45: 97 – 99.
- Richardson, J. E., Fay, M. F., Cronk, Q. C. B., Bowman, D. & Chase, M. W. (2000). A phylogenetic analysis of Rhamnaceae using *rbcL* and *trnL-F* plastid DNA sequences. *Amer. J. Bot.* 87: 1309 – 1324.
- Rohlf, F. J. (1998). NTSYS-pc numerical taxonomy and multivariate analysis system version 2.0. Applied Biostatistics INC., New York.
- Roldàn-Ruiz, I., Dendauw, J., Van Bockstaele, E., Depicker, A. & De Loose, M. (2000). AFLP markers reveal high polymorphic rates in ryegrasses (*Lolium* spp.). *Molec. Breed.* 6: 125 – 134.
- Rowe, N. & Speck, T. (2005). Plant growth forms: an ecological and evolutionary perspective. *New Phytol.* 166: 61 – 72.
- Savelkoul, P. H. M., Aarts, H. J. M., De Haas, J., Dijkshoorn, L., Duim, B., Otsen, M., Rademaker, J. L.W., Schouls, L. & Lenstra, J. A. (1999). Amplifiedfragment length polymorphism analysis: The state of an art. J. Clin. Microbiol. 37: 3083 – 3091.
- Suraninpong, P., Nuanlaong, S. & Wuthisuthimethavee, S. (2015). A new classification of Thailand's *Nepenthes* species by genetic analysis of AFLP markers. *Acta Hortic.* 1100: 77 – 82.
- Tamizi, A. A., Ghazalli, M. N., Nikong, D., Besi, E. E., Mat-Esa, M. I., Mohd-Nordin, A. R., Latiff, A. & Shakri, M. A. (2020). *Nepenthes malayensis* (Nepenthaceae), a new species of carnivorous pitcher plant from Peninsular Malaysia. *Kew Bull.* 75: 63.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., Lee, T. V. D., Hornes, M., Frijters, A., Pot, J., Paleman, J., Kuiper, M. & Zabeau, M. (1995). AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res.* 23: 4407 – 4414.

#### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.