

utrinque 3-5 subtus prominentibus, pedunculis axillaribus quam folia brevioribus gracilibus unifariam puberulis apice 1-vel 3-floris bracteatis, bracteis subulatis glabris, pedicellis brevibus patentibus vel subdeflexis glabris, calyce campanulato ad medium quinquifido glabro dentibus deltoideo-attenuatis acutis, corollæ tubo subrecto sub-compresso calyce duplo longiore, limbo inæqualiter 5-lobo cœruleo lobo inferiore cuneato-ovovato subtruncato lobis intermediis brevioribus ellipticis obtusis superioribus majoribus oblique ellipticis obtusis, staminibus styloque longe exsertis incurvis, fructu quadrilobo glabro.

*Habitat*.—South Africa: Natal, Gerrard, 1252, Mooi River Valley, 2000-3000 ft., Sutherland; Swaziland, Mrs. K. Saunders.

*Poli* 1-6 lin. longi, laminæ 4 lin.- $2\frac{1}{4}$  poll. longæ, 2 lin.-1 poll. late. *Pedunculi*  $\frac{3}{4}$ - $1\frac{1}{4}$  poll. longi. *Bracteæ* 1- $1\frac{1}{2}$  lin. longæ. *Pedicelli* 1-2 lin. longi. *Corollæ* tubus  $3\frac{1}{2}$  lin. longus, limbus 7-8 lin. diam. *Stamina* 8 lin. longa.

Allied to *C. myricoides*, R. Br., but easily recognised by its much narrower and very acute calyx-teeth.

178. *Clerodendron polycephalum*, Baker [Verbenaceæ]; fruticosum, erectum, ramulis validis dense ferrugineo-pilosus, foliis ternatis vel oppositis distincte petiolatis evatis cordatis cuspidatis, integris utrinque viridibus pubescensibus, floribus in cimas multifloras conglomeratas pedunculatas terminales dispositis, pedunculis pedicellisque pubescensibus, calyce piloso tubo obconico dentibus ovatis tubo æquifongis, corollæ tubo cylindrico calyce 2-3-plo longiore, lobis parvis obovatis, staminibus elongatis.

*Habitat*.—Interior of western Lagos, Dr. Rowland.

*Folia* 3-4 poll. longa, 2- $2\frac{1}{2}$  poll. lata. *Calyx*  $1\frac{1}{2}$  lin. longus. *Corollæ* tubus 3-4 lin. longus, limbus expansus  $1\frac{1}{2}$  lin. diam.

Ranks amongst the small-flowered species of the subgenus *Euclerodendron* near *C. formicarum*, Gürke in Engl. Jahrb. XVIII. 179.

179. *Nepenthes Smilesii*, Hemsl. [Nepenthaceæ]; parva, acaulescens? vel saltem ramulis floriferis interdum valde abbreviatis fere undique plus minusve puberulis, foliis confertis angustis utrinque attenuatis, ascidiis mediocribus recurvo-erectis fere cylindricis, costis anticis anguste alatis fimbriato-ciliatis vel fere obsoletis, calcare obsoleto, peristomio cylindraceo transversim crebre costato, operculo fere orbiculari infra multiglandulos, floribus ♂ ferrugineo-pubescentibus simpliciter racemosis, racemis longe pedunculatis, pedicellis brevissimis, perianthii segmentis ovalibus concavis, antheris 8 uniseriatis.

*Habitat*.—Northern Siam: grows on grass land on the ground, at Baw Saw, Nam Kawng, F. H. Smiles.

*Lamina* folii 4-6 poll. longa. *Cirrus* 1-2 poll. longus. *Ascidia* 2-3 poll. longa. *Operculum* 9-15 lin. latum. *Scapus* 10-12 poll. altus.

Mr. Smiles describes this as having a green pitcher with a red cover and a red border.

180. *Ophiopogon clavatus*, Wright [Hæmodoraceæ-Ophiopogoneæ]; herba, rhizomate gracili repente squamis sparse vestito, foliis aggregatis